

Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/SE04/001879

International filing date: 15 December 2004 (15.12.2004)

Document type: Certified copy of priority document

Document details: Country/Office: SE
Number: 0303364-4
Filing date: 15 December 2003 (15.12.2003)

Date of receipt at the International Bureau: 03 January 2005 (03.01.2005)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse

PRV

PATENT- OCH REGISTRERINGSVERKET

Patentavdelningen

PCT / SE 2004 / 001879

Intyg Certificate

Härmed intygas att bifogade kopior överensstämmer med de handlingar som ursprungligen ingivits till Patent- och registreringsverket i nedannämnda ansökan.

This is to certify that the annexed is a true copy of the documents as originally filed with the Patent- and Registration Office in connection with the following patent application.




(71) Sökande Nitricare HB, Uppsala SE
Applicant (s)

(21) Patentansökningsnummer 0303364-4
Patent application number

(86) Ingivningsdatum 2003-12-15
Date of filing

Stockholm, 2004-12-21

För Patent- och registreringsverket
For the Patent- and Registration Office


Görel Gustafsson

Avgift
Fee

New device and method for prevention of nosocomial infections

The present invention concerns a device and method for reducing the incidence of nosocomial infections in humans and animals having an invasive medical device inserted into the body, in particular catheter-associated urinary tract infections. More specifically, the invention relates to a device having means for releasing low molecular antimicrobial compounds that permeates to the adjacent tissue and/or body cavity.

Background

Nosocomial infections are infections that are caught during a treatment or procedure performed in a hospital. Nosocomial infections are a major global concern that leads to increased hospitalisation, and sometimes even permanent debilities for the patient. In addition to the consequences for the patient, nosocomial infections may infect other patients and cause increased health care and hospital costs.

Major sources of nosocomial infections are insufficient sterilisation of medical equipment and unsatisfactory hygiene of the personnel at hospitals and other care centres and nursing homes. Outbreaks of nosocomial infection have been linked to a variety of non-sterile equipment like electronic thermometers and blood pressure cuffs.

Staphylococcus aureus is the most common cause of nosocomial infections and is of increasing concern because of the spread of methicillin resistant strains, which are refractory to treatment by most antibiotics. Examples of other microbial pathogens causing nosocomial infections are *Escherichia coli*, *Serratia marcescens*, *Klebsiella* and *Enterobacter* species and *Candida* species.

Venous catheters, urinary catheters, and ventilator-associated tubings are common sites and causes of nosocomial infections. Actually, catheter-associated urinary tract infection is the most common nosocomial infection and is, together with nosocomial pneumonia, two of the major causes of hospital-acquired infection, for which a substantial proportion of prescribed antibiotics is used.

Urinary catheterisation is a routine procedure in the hospital and chronic care settings and is associated with a significant risk of infection. The incidence of catheter-associated urinary tract infection (CAUTI) approaches 50 % after three days of continuous catheterization. The presence of a catheter within the urinary tract may also increase the difficulty of treating the infection. If a urinary catheter is left in place for long periods of time, bacteria will inevitably grow in it. A harmful infection may occur if the number of bacteria

becomes large or if specific pathologic bacteria grow in the urinary tract. Complications of CAUTI include bacteremia, pyelonephritis, urinary stones and renal failure with resulting morbidity and increased risk of death.

Most microorganisms, except *Staphylococcus aureus*, causing endemic CAUTI
5 derive from the patient's own colonic and perineal flora or from the hands of health-care personnel during catheter insertion or manipulation of the collection system. Extraluminal contamination may occur early when the catheter is inserted or later by microorganisms ascending from the perineum by capillary action in the thin mucous film close to the external catheter surface. Intraluminal contamination occurs by reflux of microorganisms gaining
10 access to the catheter lumen due to breakage in closed drainage or contamination of collection bag urine.

Most urinary catheters are equipped with an inflatable cuff situated near the catheter tip. The cuff keeps the catheter in place in the bladder after insertion. When inflated it represents the largest surface area of the catheter in the bladder. The outflow orifice of the
15 catheter is situated distal to the cuff meaning that the cuff itself will be constantly embedded in residual urine. It is well known that urine is an excellent growth medium for urinary pathogens. Consequently, the cuff is a particularly vulnerable site for bacterial adherence and growth.

There is a heavy use of systemic antimicrobial drugs to treat and prevent
20 nosocomial infections. Antimicrobial drugs probably keeps the rate of hospital-acquired infections at a considerably lower rate than it would be otherwise, but it unfortunately selects for resistant microorganisms causing many of the most severe nosocomial infections. Thus, there is a great need for new methods to prevent the occurrence and spread of nosocomial infections, both in an economical perspective but also from the patient's point of view.

25

Prior art

Most infected urinary catheters are covered by a thick biofilm containing the infecting microorganisms embedded in a matrix of host proteins and microbial exoglycocalyx. Biofilms may appear both intraluminally and extraluminally. Anti-infective-impregnated and
30 silver-hydrogel catheters, which inhibit adherence of microorganisms to the catheter surface, significantly reduce the risk of CAUTI. However, those coated catheters mainly affect the infections caused by grampositive organisms or yeasts adhering to the surface of the catheter. Silver alloy and silver oxide coated catheters are also widely used to inhibit establishment of infection.

Other attempts to inhibit CAUTI are the use of anti-infective lubricants when inserting the catheter, soaking the catheter in anti-infective solution before insertion and continuously irrigating the catheterised bladder with anti-infective solution. Efforts have also been made to seal the connection between the catheter and collection tubing. None of these
5 methods have given satisfactory results and new better ways to inhibit nosocomial infections would be valuable to replace or complement existing methods.

Nitric oxide (NO) is known for its role in the defence against microorganisms. US Patent Application No. 2002/0155174 describes the use of acidified nitrite as an antimicrobial agent for the treatment of viral diseases of the skin by topical application
10 thereto. Acidified nitrite forms nitrous acid, which in turn dissociates to form oxides of nitrogen.

Nablo and Schoenfisch (J Biomed Mater Res, 67A: 1276-1283, 2003) also utilises the antibacterial properties of nitric oxide. They describe NO-releasing sol-gel coatings decreasing bacterial adhesion by 30 to 95% that, for example, can be used on
15 implanted medical devices to prevent biofilm formation.

None of the above mentioned methods for preventing CAUTI relates to the cuff and preventing biofilm formation and bacterial growth on the surface of the cuff. For example, in silver coated catheters the inflatable cuff is not possible to coat and will remain a particularly susceptible site for infection.

20 In the present invention antimicrobial low molecular compounds are used to achieve an effective means to prevent establishment of nosocomial infections, in particular CAUTI.

Summary of the invention

25 The present invention relates to a device for insertion in a human or animal body and/or body cavity, said device having inflatable/expandable means for securing said device. The inflatable/expandable means comprises at least one component capable of releasing at least one low molecular antimicrobial compound (LMAC) in the device, said LMAC permeating to the adjacent tissue and/or body cavity.

30 The present invention also relates to a method for the prevention and treatment of nosocomial infections, such as CAUTI, originating from the insertion and/or use of a medical device in a human or animal. The method is characterized in that at least one component capable of releasing at least one LMAC is present in an inflatable/expandable

means of the device or administered into the device. The LMAC is capable of permeating to the adjacent tissue and/or body cavity.

Short description of the drawings

5 The present invention will be described in closer detail in the following description, examples, and attached drawings, in which

Fig. 1 schematically shows a double lumen urinary catheter having a hollow elongated body 1, an inlet/outlet 2 with a suitable fitting, and an inlet/outlet 3, also with a suitable fitting. The tip of the catheter has an opening 4, which is in fluid connection through
10 a first lumen with the inlet/outlet 2. Slightly below the tip, a cuff 5 is provided. The cuff can be inflated through an opening 6, in fluid connection through a second lumen with the inlet/outlet 3.

Fig. 2 schematically shows the end part of any device 7 to be introduced into the body or a body cavity, and having a first lumen 8 for administering and a second lumen 9 for
15 inflating / deflating a cuff delimited by an elastic membrane 10. The volume of the inflated cuff is indicated as 11.

Fig. 3 schematically shows a system for urine collection where a catheter 12 is inserted in a bladder 13. The residual urine volume is indicated as 14. The catheter is connected via connecting means 15 to a collection bag 17, containing a volume of urine
20 indicated as 17. The capital letters A, B, C, and D indicated critical points of contamination and/or bacterial growth.

Fig. 4 shows the experimental set-up described in the example, where a sample of *E. coli* infected urine was placed in a 50 ml flask 19. An all-silicone catheter 18 was inserted in the flask, the cuff inflated with a solution according to one embodiment of the
25 invention, and the flask turned up-side-down. The cuff efficiently sealed the neck of the flask and a volume of urine 20 was trapped in the flask.

Fig. 5 shows schematically two embodiments where the catheter tip is modified so that the LMAC will penetrate also to the inner lumen of the catheter and thus exert its antimicrobial effect on the urine collected from the bladder. In A the lumen wall has local
30 areas of reduced thickness, here shown as wells in the catheter wall. In B the catheter wall is thinner at the section covered by the cuff, or part of said section.

Detailed description of the invention

Whenever a medical device comes in contact with a patient, a risk of infection is created. The risk of infection dramatically increases for invasive medical devices. The present invention provides a device and method for reducing the risk of nosocomial infections and for the treatment of nosocomial infections. Nosocomial infections may arise both inside and on the outside of an invasive medical device. Thus, the simultaneous reduction and elimination of microbial organisms both on the inside and outside of the device would be beneficial.

The present inventors surprisingly found that the use of low molecular antimicrobial compounds (LMAC's) could solve this problem. More specifically, the present invention relates to a device for insertion in a human or animal body and/or body cavity, said device having inflatable/expandable means for securing the device (Fig. 1 and 2), wherein the inflatable/expandable means comprise at least one component capable of releasing at least one LMAC in the device, said LMAC permeating to the adjacent tissue and/or body cavity.

The device of the present invention is an invasive medical device. An invasive medical device refers to any device wherein at least a portion of the device may be inserted percutaneously or into any site of the body of a human or an animal. Examples of medical devices includes urinary catheters, intratracheal tubes, vascular catheters, vascular catheter ports, wound drain tubes etc. In particular the present invention relates to a catheter for insertion into the urinary tract or an intratracheal tube for insertion into the respiratory tract of a human or animal body.

The LMAC released in the securing means of the device is a compound capable of permeating said means to the adjacent tissue and/or body cavity thereby exerting its antimicrobial effect both on the inside and on the outside of the device. Thus, the LMAC can be any antimicrobial compound, such as an antimicrobial gas, with sufficiently low molecular weight to permeate the device. In particular the LMAC is a reactive nitrogen intermediate (RNI), a reactive oxygen intermediate (ROI) or a combination of these two. Examples of LMAC's are, but not limited to, nitric oxide (NO), NO₂, N₂O₃, N₂O₄, HNO₂, HNO₃, NO⁺, NO⁻, O₂⁻, O₃, singlet oxygen, H₂O₂, OONO⁻, HOONO, OH radical and HOCl.

The inflatable/expandable means of the device can be any means for securing a medical device inserted into a human or animal body or body cavity. In the present invention the inflatable/expandable means consist of an elastic material of sufficient medical grade being permeable to gas but impermeable to water and which is expandable and inflatable to secure the device in the body. In particular the inflatable/expandable means is a cuff consisting of, but not limited to, elastic materials comprising polysiloxanes such as silicone

rubber comprising polydimethylsiloxanes, latex-free rubber, silicon-coated rubber or a semi-permeable or selectively permeable membrane such as Goretex®.

The LMAC's exert their antimicrobial effect unspecifically, which probably reduces the potential of the microorganisms to develop resistance to the LMAC's. The use of LMAC's with short half-lives in relatively high dose will further reduce the development of resistant bacteria. Furthermore, since some RNIs and ROIs are generated endogenously, the use of such compounds is unlikely to constitute any major health risk to the patient. Another advantage with the present invention is the local production of antibacterial compounds, which will not affect the normal microbial flora of, for example, the gut. The use of antibiotics in the inflatable/expandable means cannot exert the same antimicrobial effect as LMAC's since antibiotics can not permeate said inflatable/expandable means. Furthermore, the LMAC's permeating the said inflatable/expandable means will lead to high local concentrations of antimicrobial compounds leading to efficient biofilm prevention.

The LMAC of the present invention is released from at least one component comprised in the inflatable/expandable means. Said at least one component releasing the LMAC is any compound capable of releasing an antimicrobial compound with sufficiently low molecular weight to permeate the device. Examples of components that release RNIs/ROIs include, but are not limited to, S-nitrosothiols (low or high molecular-weight thiols), NONOates, nitroprusside, organic nitrate/nitrites, N-nitroso compounds, C-nitroso compounds, oximes, sydnonimines, oxadiazoles (furoxans), oxatriazoles, nitroxyl generating compounds (e.g. pyruvic acid), hydroxylamine, N-hydroxy-guanidines, nitrosylchloride, sodium azide, nitrosylhydrogensulphate, nitrosyltetrafluoroborate, dinitrosyl-iron-cysteine complex, etc.

There are several factors affecting the release of the LMAC. The RNIs/ROIs released by those compounds have different half-lives and differ in their pH dependency. Thus, the at least one component releasing the LMAC can be alternated depending on the special circumstances associated with the infection, the patient, where in the body the device is inserted etc.

Different LMAC's exert diverse effects on different bacteria. Thus, the present invention gives the possibility of using different LMAC's depending on the infectious bacteria. For example, *S. aureus* is sensitive to NO alone while *E. coli* is not. Furthermore, the LMAC's released can also be changed or modified with respect to the localization of the device since, for example, bacteria causing nosocomial infections coupled to the use of a urinary catheter might, at least partly, be distinct from the bacteria causing infections related

to the use of an intratracheal tube. Depending on the patient and the type of infection, the dosage of released LMAC can be changed or modified through the present invention. The term "changed or modified" as used above comprises changing the LMAC itself, e.g. by substituting one LMAC for another or using different combinations of LMAC's, or by
5 changing the concentration(s) of the same.

In one embodiment of the present invention the LMAC is released when said at least one component is contacted with a second component. The second component can, for example, be an acid or base initiating the release of the LMAC when contacted with the component releasing the LMAC.

10 In another embodiment the component releasing the LMAC and the second component are present in the securing means as dry powders, granulaes, thin films or the like coating the inside of the cuff and the contact between the two components is accomplished through the addition of a liquid to said means. The liquid can for example be, but is not limited to, of water, saline or any physiological buffer, such as phosphate buffered saline
15 (PBS) or the like.

In one embodiment of the present invention the at least one component of the device releasing the LMAC releases the LMAC upon acidification. The acidification is achieved through the addition of an acidifying agent (the "second component") reducing the pH in the device to about pH 1 - 5.5, preferably to about 2 - 4. The acidifying agent can be
20 any acid reducing the pH in the device to the above-mentioned pH. Appropriate acids are for example, but not limited to, ascorbic acid, citric acid, acetic acid, butyric acid, diluted hydrochloric acid, diluted sulphuric acid, formic acid, nitrous acid or nitric acid.

In one embodiment of the present invention said at least one component is nitrite and said second component is an acidifying agent that, when brought into contact, release
25 RNIs such as nitric oxide, N_2O_3 and nitrous acid.

In one embodiment of the present invention the means for securing the inventive device is at least one inflatable cuff situated somewhere along the device and the at least one component releasing the LMAC is present in said inflatable cuff. In another embodiment the second component is present in said inflatable cuff. In yet another embodiment both said at
30 least one component releasing a LMAC and the second component are present in said inflatable cuff. The inflatable cuff has an arrangement for introducing said at least one component or said second component in the cuff. Said arrangement of the inflatable cuff making it possible to administer the component(s) repeatedly during the use of the device, thus more efficiently preventing or treating nosocomial infections during prolonged use of the

device in a patient. Said arrangement is also suitable for the introduction of a liquid into the cuff contacting the at least one component releasing a LMAC and the second component.

When the device is a catheter (Fig. 3) for insertion into the urinary tract of the human or animal body, the cuff, when the catheter is inserted, is situated in the urinary bladder surrounded by the residual urine present in the bladder. Thus, the LMAC's released by the present invention solves the problem with infections establishing on the outside of the cuff surrounded by the residual urine.

In one embodiment of the present invention the part of the device surrounded by the inflatable/expandable means have local areas of reduced thickness in order to allow the LMAC's to permeate also to the inside of the device thus exerting an antimicrobial effect also inside the device (Fig. 5A). According to another embodiment the catheter wall is thinner at a section surrounded by the inflatable cuff or a part thereof (Fig. 5B).

The present invention also relates to a method for preventing and/or treating nosocomial infections originating from the insertion and/or use of a device inserted into a human or animal body and/or body cavity, said device having inflatable/expandable means for securing the device, wherein at least one component capable of releasing at least one LMAC in said means is administered into said means, said LMAC permeates to the adjacent tissue and/or body cavity. In one embodiment of the present invention the method is used for reducing the risk of and treating nosocomial infections originating from the insertion and use of a urinary catheter. In another embodiment the method is used for reducing the risk of and treating nosocomial infections originating from the insertion and use of an intratracheal tube inserted into the respiratory tract of a human or animal body.

The inflatable/expandable means of the device can be any means for securing a medical device inserted into a human or animal body or body cavity. In the inventive method the inflatable/expandable means consists of an elastic material of sufficient medical grade being permeable to gas but impermeable to water and which is expandable and inflatable to secure the device in the body. In particular the inflatable/expandable means is a cuff consisting of, but not limited to, an elastic material consisting of or comprising polysiloxanes such as silicone rubber comprising polydimethylsiloxanes, latex-free rubber, silicon-coated rubber or a semi-permeable or selectively permeable membrane such as Goretex®. When the device is a urinary catheter the cuff, when inserted into the human or animal body, is situated in the urinary bladder surrounded by the residual urine present in the bladder.

The low molecular compound exerting the antimicrobial effect can be any antimicrobial compound with sufficiently low molecular weight to permeate the

inflatable/expandable means of the device wherein it is released. In particular the LMAC is a RNI, a ROI or a combination of these two. Examples of LMAC's are, but not limited to, nitric oxide (NO), NO₂, N₂O₃, N₂O₄, HNO₃, HNO₂, NO⁺, NO⁻, O₂⁻, O₃, singlet oxygen, H₂O₂, OONO⁻, HOONO, OH radical and HOCl.

5 The LMAC of the inventive method is released from at least one component comprised in the securing means of the device. Said at least one component releasing the LMAC is any compound capable of releasing an antimicrobial compound with sufficiently low molecular weight to permeate the device. Examples of components that release RNIs/ROIs include, but are not limited to, S-nitrosothiols (low or high molecular-weight
10 thiols), NONOates, nitroprusside, organic nitrate/nitrites, N-nitroso compounds, C-nitroso compounds, oximes, sydnonimines, oxadiazoles (furoxans), oxatriazoles, nitroxyl generating compounds (e.g. pilotys acid), hydroxylamine, N-hydroxy-guanidines, nitrosylchloride, sodium azide, nitrosylhydrogensulphate, nitrosyltetrafluoroborat, dinitrosyl-iron-cysteine complex.

15 In one embodiment of the method according to the invention the LMAC is released when said at least one component is contacted with a second component. The second component can, for example, be an acid or base initiating the release of the LMAC when contacted with the component releasing the LMAC.

 In another embodiment of the method the component releasing the LMAC and
20 the second component are present in the device as dry powders, granulaes, thin films or the like and the contact between the two components is accomplished through the addition of a liquid to the device. The liquid can for example be, but is not limited to, water, saline or any physiological buffer, such as phosphate buffered saline (PBS) or the like.

 In yet another embodiment of the inventive method the at least one component
25 releasing the LMAC releases the low molecular compound upon acidification. The acidification is achieved through the addition of an acidifying agent reducing the pH in the device to about pH 1 - 5.5, preferably to about 2 - 4. The acidifying agent can be any acid reducing the pH in the device to the above-mentioned pH. Appropriate acids are for example ascorbic acid, citric acid, acetic acid, butyric acid, diluted hydrochloric acid, diluted sulphuric
30 acid, formic acid, nitrous acid, or nitric acid.

 In another embodiment of the inventive method said at least one component is nitrite and said second component is an acidifying agent that, when brought into contact, release RNIs such as nitric oxide, N₂O₃ and nitrous acid.

In another embodiment of the present invention the method is for treatment of nosocomial infections in and around an inflatable cuff situated along the device. The at least one component and/or the second component is/are administered into the cuff situated along the device. The inflatable cuff has an arrangement (3 and 6 in Fig. 1) making it possible to
5 administer said component that releases at least one LMAC and/or said second component to the cuff. Through the administrative arrangement of the inflatable cuff said component releasing a low molecular compound and/or the second component can be added to the device repeatedly during the use of the device, thus more efficiently preventing or treating nosocomial infections during prolonged use of the device in a patient. Said arrangement is
10 also suitable for the administration of a liquid into the cuff contacting the at least one component releasing a LMAC and the second component. Furthermore, through the use of said arrangement it is easy to change or modify the dosage of the released LMAC in each individual patient.

The invention also relates to a method for the prevention and treatment of
15 nosocomial infections, wherein a device as described above is used.

Urinary collection devices, such as containers or drainage bags, are generally used for collecting urine from a catheterized patient. A collection device usually has tubings that are attached to the catheter leading the urine to the collection device (Fig. 3). Catheterization often results in the possibility of urinary tract infection resulting from growth of
20 microorganisms in the collection device and the associated tubing. The microorganisms growing in the collection device might be transported back to the bladder when the collection system is moved or manipulated in association with breakage in closed drainage. There are four main critical points for establishment of infection during catheterization as indicated in Fig. 3. One is the area surrounding the opening of the urinary tract, indicated as A. Another is
25 the inside of the bladder, B, where the cuff and the tip of the catheter reside in residual urine. A third point of infection is the connection or coupling C between the catheter itself and the collection device. In Fig. 3 the coupling C is drawn as situated closer to the catheter than to the collection bag. In reality, C could be at the collection bag, any where between the catheter and said bag, or in fact, several couplings C could be provided. The fourth critical point is the
30 collection bag itself, D, where bacteria can grow in large volumes of urine at room temperature. There is a need for an effective method for the prevention of urinary tract infection originating from contaminants in the collection system.

Thus, the present invention also relates to a method for prevention of nosocomial infections in patients having a urinary catheter by preventing microbial growth in

the collection device coupled to a catheter wherein at least one component releasing a LMAC is added to the collection device. An advantage with the use of the inventive method is that the use of LMAC, exerting its antimicrobial effect unspecifically, most likely will have lower probability of giving rise to antibiotic resistant microbial strains. In particular the LMAC is a RNI, a ROI or a combination of these two. Examples of LMAC's are, but not limited to, nitric oxide (NO), NO₂, N₂O₃, N₂O₄, HNO₃, HNO₂, NO⁺, NO⁻, O₂⁻, O₃, singlet oxygen, H₂O₂, OONO⁻, HOONO, OH radical and HOCl.

The at least one component releasing the LMAC in the collection device is any compound capable of releasing a LMAC in the collection device, such as, but not limited to, S-nitrosothiols (low or high molecular-weight thiols), NONOates, nitroprusside, organic nitrate/nitrites, N-nitroso compounds, C-nitroso compounds, oximes, sydnonimines, oxadiazoles (furoxans), oxatriazoles, nitroxyl generating compounds (e.g Piloty's acid), hydroxylamine, N-hydroxy-guanidines, nitrosylchloride, sodium azide, nitrosylhydrogensulphate, nitrosyltetrafluoroborat, dinitrosyl-iron-cysteine complex.

In another embodiment of the method the at least one component releasing the LMAC in the collection device releases the low molecular compound upon acidification. The acidification is achieved through the addition of an acidifying agent ("the second component") reducing the pH in the device to about pH 1 - 5.5, preferably to about 2 - 4. The acidifying agent can be any acid reducing the pH in the device to the above-mentioned pH.

Appropriate acids are for example ascorbic acid, citric acid, acetic acid, butyric acid, diluted hydrochloric acid, diluted sulphuric acid, formic acid, nitrous acid, or nitric acid.

In another embodiment of the inventive method said at least one component is nitrite and said second component is an acidifying agent that, when brought into contact, release RNIs such as nitric oxide, N_2O_3 and nitrous acid in the collection device.

The present invention also relates to a kit to be used for preventing or treating nosocomial infections originating from the insertion and/or use of an invasive medical device, such as a urinary catheter or intratracheal tube, having an inflatable cuff in a human or animal body and/or body cavity. Said kit comprises said device and a syringe suitable for inflating said cuff and said syringe comprises the necessary components for the release of a LMAC after administration of said components into said inflatable cuff of said device. The LMAC can be any compound with sufficiently low molecular weight to permeate the cuff and/or the device. In one embodiment the LMAC is a reactive nitrogen intermediate (RNI), a reactive oxygen intermediate (ROI) or a combination of these two. The kit is easy to use and there is

no need for additional equipment or special training of the personnel at hospitals or other health care centres.

The necessary components comprised in the syringe are at least one component releasing the LMAC and, optionally, a second component inducing the release of the LMAC upon contact with the LMAC releasing component. The component(s) can be present as dry
5 powders, granulaes, thin films or the like that release said LMAC upon combination with a liquid such as water, saline or any physiological buffer, such as PBS. In one embodiment the powders and the liquid are kept in separate containers until combined in the syringe prior to administration.

10 In one embodiment of the inventive kit one of the necessary components in the syringe is nitrite that upon acidification produces RNIs. The acidification can for example be achieved through the addition of, but not limited to, ascorbic acid, citric acid, acetic acid, butyric acid, diluted hydrochloric acid, diluted sulphuric acid, formic acid, nitrous acid, or nitric acid.

15 The present invention can also be used in combination with other devices, methods and/or kits developed for the prevention and/or treatment of nosocomial infections, such as, but not limited to, antiinfective lubricants that can be used when inserting the medical device, antiinfective creams or ointments applied to the device, soaking of the medical device in antiinfective antimicrobial-drug solution prior to use, silver oxide coated
20 devices (e.g. catheters).

The present invention also relates to a device as described above exerting its antimicrobial effect on non-nosocomial infections. In one embodiment the at least one component releasing at least one LMAC is present in an inflatable cuff situated along a catheter inserted in the urinary tract of a human. The LMAC penetrates the cuff and exert its
25 effect on the prostate. It has been suggested that for example chronic prostatitis is caused by a bacterial infection. By arranging a second cuff at a specific distance from the first cuff, securing the device, the second cuff can be positioned in the urinary tract so that it is surrounded by the prostate gland. This embodiment makes possible a long term, minimally invasive treatment of the prostate.

30 The present invention will now be further described in the following non-limiting example.

Example

Materials & Methods

An *E. coli* strain isolated from urine of a patient with lower urinary tract infection was used. A 50 ml glass bottle with a narrow neck (Fig. 4) was filled with fresh
5 urine (pH 6.5) from a healthy volunteer. The urine was inoculated with the *E. coli* strain to a final density of 10^5 colony forming units/ml. Then an all-silicone urinary catheter was inserted in the bottle and the cuff was filled with a solution comprising:

1. Saline + ascorbate (20 mM) + HCl to a final pH of 2.0 (control; n=3)
10 or
2. Saline + ascorbate (20 mM) + HCl + sodium nitrite (2 mM), pH 2 (nitrite; n=3)

The expanded cuff was fixed at the neck of the bottle to prevent leakage of urine when the bottle was turned up-side-down. The glass bottle was then incubated in 37° C for 10
15 h after which growth of *E. coli* in the surrounding urine was monitored by optical density (OD) at 540 nm on a Spectramax® (Molecular Devices Inc.).

Results

In controls OD values increased from 0.12 to 0.35 while in the nitrite group, no
20 visible growth was observed (OD 0.12 before and 0.13 after 10 h).

The experiment shows that the addition of nitrite to an acidic ascorbate solution results in generation of bacteriostatic compounds (such as NO, N₂O₃ and nitrous acid), which can penetrate a thin silicone membrane and inhibit bacteria growing in urine outside the
membrane.

25

Although the invention has been described with regard to its preferred
embodiments, which constitute the best mode presently known to the inventors, it should be
understood that various changes and modifications as would be obvious to one having the
ordinary skill in this art may be made without departing from the scope of the invention which
30 is set forth in the claims appended hereto.

Claims

1. A device for insertion in a human or animal body and/or a body cavity, said device having inflatable/expandable means for securing said device, **characterized in** that said inflatable/expandable means comprise at least one component capable of releasing at least one low molecular antimicrobial compound (LMAC) in the device, said LMAC permeating to the adjacent tissue and/or body cavity.
2. A device according to claim 1, wherein the LMAC is released when said at least one component is contacted with a second component.
3. A device according to claim 2, wherein said contact is accomplished through the introduction of a liquid selected from the group consisting of water, saline or any physiological buffer to said means.
4. The device according to claim 1, wherein said device is a catheter for insertion into the urinary tract of said human or animal and said means for securing said device is an inflatable cuff.
5. The device according to claim 4, wherein said cuff when inserted into the urinary tract is situated in the urinary bladder.
6. The device according to claim 1, wherein said device is an intratracheal tube.
7. The device according to claim 1, wherein said LMAC is a reactive nitrogen intermediate, a reactive oxygen intermediate or a combination of these two.
8. The device according to claim 1, wherein the LMAC is selected from the group consisting of nitric oxide (NO), NO₂, N₂O₃, N₂O₄, HNO₃, HNO₂, NO⁺, NO⁻, O₂⁻, O₃, singlet oxygen, H₂O₂, OONO⁻, HOONO, OH radical and HOCl.
9. The device according to claim 1, wherein said at least one component releasing a LMAC releases the LMAC upon acidification.
10. The device according to claim 9, wherein said at least one component is nitrite.

11. The device according to claim 1, wherein said at least one component releasing a LMAC releases the LMAC upon basification
- 5 12. A method for preventing and/or treating nosocomial infections originating from the insertion and/or use of a device inserted into a human or animal body and/or a body cavity, said device having inflatable/expandable means for securing said device, **characterized in** that at least one component capable of releasing at least one low molecular antimicrobial compound (LMAC) in said means is administered to said
10 means, said LMAC permeating to the adjacent tissue and/or body cavity.
13. The method according to claim 12, wherein the LMAC is released when said at least one component is contacted with a second component.
- 15 14. The method according to claim 13, wherein said contact is accomplished through the introduction of a liquid selected from the group consisting of water, saline or any physiological buffer to said means.
- 20 15. The method according to claim 13 or 14, wherein said at least one component that releases a LMAC, said acidifying agent and/or said liquid is/are administered into said means for securing the device, said means having an arrangement for administering said component that release at least one low molecular antimicrobial compound, said acidifying agent in the cuff and/or said liquid.
- 25 16. The method according to claim 12, wherein said device is a catheter for insertion into the urinary tract of said human or animal body and said means for securing the device is an inflatable cuff.
- 30 17. The method according to claim 12, wherein said device is an intratracheal tube for insertion into the respiratory tract of said human or animal body.
18. The method according to claim 12, wherein said LMAC is a reactive nitrogen intermediate, a reactive oxygen intermediate or a combination of these two.

19. The method according to claim 12, wherein the LMAC is selected from the group consisting of nitric oxide (NO), NO₂, N₂O₃, N₂O₄, HNO₃, HNO₂, NO⁺, NO⁻, O₂⁻, O₃, singlet oxygen, H₂O₂, OONO⁻, HOONO, OH radical and HOCl.
- 5 20. The method according to claim 12, wherein said at least one component releases the LMAC upon acidification.
21. The method according to claim 20, wherein said at least one component is nitrite.
- 10 22. The method according to claim 12, wherein said at least one component releases the LMAC upon basification.
23. A method for the prevention and treatment of nosocomial infections, characterized in that a device according to any one of the claims 1-11 is used.
- 15 24. A method for the prevention of nosocomial infections in patients having an urinary catheter inserted by preventing microbial growth in the collection device coupled to an urinary catheter, **characterized in** that at least one component releasing at least one low molecular antimicrobial compound (LMAC) is added to the collection device.
- 20 25. The method according to claim 24, wherein said at least one component is nitrite that upon acidification releases a LMAC in the collection device.
- 25 26. The method according to claim 25, wherein the acidification is accomplished through the addition of an acidifying agent reducing the pH in the device to about pH 1 - 5.5, preferably to about pH 2 - 4.
- 30 27. A kit to be used in the prevention and/or treatment of nosocomial infections originating from the insertion and/or use of a invasive medical device having an inflatable cuff in a human or animal body and/or body cavity, said kit comprising said device and a syringe suitable for inflating said cuff, said syringe comprising the necessary components for the release of at least one low molecular antimicrobial compound (LMAC) after administration of said components into said inflatable cuff of said device.

28. A kit according to claim 27, wherein the device is a urinary catheter.

29. A kit according to claim 27, wherein the device is an intratracheal tube.

5

30. The kit according to claim 27, wherein said necessary components are present as powders that release said LMAC upon combination with a liquid such as water, saline or any physiological buffer.

10

31. The kit according to claim 27, wherein said necessary components are present as separate solutions that are combined prior to administration or simultaneously with the inflation of said cuff.

15

1-1-15

Abstract

The present invention discloses a device and a method for reducing the risk of hospital-acquired infections, nosocomial infections, originating from the insertion of the device and/or use of the device in the body of a human or animal. The invention in particular
5 relates to a device releasing at least one low molecular antimicrobial compound permeating the device thereby exerting its antimicrobial effect also on the outside of the device.

(Fig. 3)

10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195